

Complications in Spine Surgery
Warwick June 2014



GLOBUS MEDICAL


Ways to Minimise Post-Operative Surgical Site Infections

Mr M J H McCarthy
Consultant Spinal Surgeon

Fusion Portfolio Minimally Invasive Surgery Biomaterials Motion Preservation

Life moves us ➤


GLOBUS MEDICAL



- Ignaz Semmelweis
- 1818-1865
- “saviour of mothers”
- Wash your hands

Life moves us ➤

GLOBUS MEDICAL



- Joseph Lister
- 1827-1912
- Sterilisation

Life moves us ➤

GLOBUS MEDICAL



- Alexander Flemming
- 1881-1955
- Penicillin

Also Louis Pasteur, Robert Koch and Ferdinand Cohn

Life moves us ➤

GLOBUS MEDICAL



2011 Outstanding Paper: Surgical Science
Morbidity and mortality of major adult spinal surgery. A prospective cohort analysis of 942 consecutive patients

John T. Street, MD, PhD^{a,*}, Brian J. Lenchan, MD^a, Christian P. DiPaola, MD^a, Michael D. Boyd, MD^a, Brian K. Kwon, MD, PhD^a, Scott J. Paquette, MD^a, Marcel F.S. Dvorak, MD^a, Y. Raja Rampersaud, MD^a, Charles G. Fisher, MD^a

- Very detailed and strict study of AEs
- 940 pts over 1 year period
- **87% had at least 1 documented complication!**
 - 39% affected LOS
 - Most common - electrolyte imbalance and medication related SEs
 - Intraop complications 10.5%
 - Post op complications 73.5%
 - 13.5% wound complication
 - **29 SSIs = 3%**
- **WHAT WE DO IS VERY HIGH RISK!**

Life moves us ➤

GLOBUS MEDICAL

Some Stats

- Spinal SSI 0.7 – 12+%
- Increased morbidity, mortality and health care costs
- SSIs = 1/3 of HAIs
- CDC Jan 2014 SSI:
 - 30 days after surgery without instrumentation
 - 1 year with instrumentation
 - Other classifications:
 - Acute (<3 weeks)
 - Subacute (3 weeks to 3 months)
 - Chronic (>3 months)

Life moves us ➤

Table 1. Criteria for Defining a Surgical Site Infection (SSI)*

Superficial Incisional SSI

Infection occurs within 30 days after the operation and infection involves only skin or subcutaneous tissue of the incision and at least one of the following:

1. Purulent drainage, with or without laboratory confirmation, from the superficial incision.
2. Organisms isolated from an aseptically obtained culture of fluid or tissue from the superficial incision.
3. At least one of the following signs or symptoms of infection: pain or tenderness, localized swelling, redness, or heat and superficial incision is deliberately opened by surgeon, unless incision is culture-negative.
4. Diagnosis of superficial incisional SSI by the surgeon or attending physician.

Do not report the following conditions as SSI:

1. Stitch abscess (minimal inflammation and discharge confined to the points of suture penetration).
2. Infection of an episiotomy or newborn circumcision site.
3. Infected burn wound.
4. Incisional SSI that extends into the fascial and muscle layers (see deep incisional SSI).

Note: Specific criteria are used for identifying infected episiotomy and circumcision sites and burn wounds.⁴¹³

Deep Incisional SSI

Infection occurs within 30 days after the operation if no implant† is left in place or within 1 year if implant is in place and the infection appears to be related to the operation and infection involves deep soft tissues (e.g., fascial and muscle layers) of the incision and at least one of the following:

1. Purulent drainage from the deep incision but not from the organ/space component of the surgical site.
2. A deep incision spontaneously dehisces or is deliberately opened by a surgeon when the patient has at least one of the following signs or symptoms: fever (>38°C), localized pain, or tenderness, unless site is culture-negative.
3. An abscess or other evidence of infection involving the deep incision is found on direct examination, during reoperation, or by histopathologic or radiologic examination.
4. Diagnosis of a deep incisional SSI by a surgeon or attending physician.

Notes:

1. Report infection that involves both superficial and deep incision sites as deep incisional SSI.
2. Report an organ/space SSI that drains through the incision as a deep incisional SSI.

Organ/space SSI

Infection occurs within 30 days after the operation if no implant† is left in place or within 1 year if implant is in place and the infection appears to be related to the operation and infection involves any part of the anatomy (e.g., organs or spaces), other than the incision, which was opened or manipulated during an operation and at least one of the following:

1. Purulent drainage from a drain that is placed through a stab wound† into the organ/space.
2. Organisms isolated from an aseptically obtained culture of fluid or tissue in the organ/space.
3. An abscess or other evidence of infection involving the organ/space that is found on direct examination, during reoperation, or by histopathologic or radiologic examination.
4. Diagnosis of an organ/space SSI by a surgeon or attending physician.

* Horan TC et al.⁴²

†National Nosocomial Infection Surveillance definition: a nonhuman-derived implantable foreign body (e.g., prosthetic heart valve, nonhuman vascular graft, mechanical heart, or hip prosthesis) that is permanently placed in a patient during surgery. If the area around a stab wound becomes infected, it is not an SSI. It is considered a skin or soft tissue infection, depending on its depth.



Jan2014


Surveillance Definitions

CDC/NHSN Surveillance Definitions for Specific Types of Infections

• **Definitions:**

- Superficial Incisional
- Deep Incisional
- Organ / Space

- Localised stitch abscess and pin site infections not classified as SSIs (common)



Life moves us ➤



CDC Jan 2014

• **All SSIs:**

- Staph aureus 30%
- Coag neg staph 13.7%
- Enterococcus 11.2%
- E.coli 9.6%
- Pseudomonas 5.6%

1. Direct inoculation at surgery
2. Contamination early post op
3. Contamination haematogenous seeding

Life moves us ➤



CDC Jan 2014

• **Endogenous:**

- Pt flora
 - Skin
 - Membranes
 - GI tract
- Seeding from distal focus infection

• **Exogenous sources:**

- Surgical personnel
 - Soiled attire
 - Aseptic tech
 - Hand hygiene
- OR environment and ventilation
- Tools, equipment, materials / implants

Life moves us ➤



Some Ps

- Patient
- Pathology (incl micro factors)
- Procedure



- Preoperative
- Perioperative
- Postoperative
- Proper
- Planning
- Prevents
- Piss
- Poor
- Performance



Life moves us ➤



Left L5/1 LRS Disc



Life moves us ➤



2009

Risk Factors for Postoperative Spinal Wound Infections After Spinal Decompression and Fusion Surgeries

Anand Veeravagu, MD,*† Chirag G. Patel, MD,*† Shivanand P. Lad, MD, PhD,*† and Maxwell Boskey, MD,*†

- 752 / 24,774 = 3% SSI
- SSI associated with:
 - Diabetes
 - Smoking
 - ASA ≥ 3
 - Preop Hct <36
 - Disseminated cancer
 - Instrumented fusion
 - Op > 3 hours

Life moves us ➤

2011

Infections in the operated spine: Update on risk management and therapeutic strategies

J.-Y. Lazennec^{a,*}, E. Fourniols^a, T. Lenoir^b, A. Aubry^c, M.-L. Pissonnier^a, B. Issartel^d, M.-A. Rousseau^a, French Spine Surgery Society

Orthopaedics & Traumatology: Surgery & Research (2011) 97S, S107–S116

- Preop hospitalization < 4/7
- Stop smoking
- Manage obesity, diabetes, malnutrition
- ID + treat infections / carriage
- Do not give prolonged post op abs
- Limit duration of tubes, lines, drains
 - Drains out <48hrs, no evidence drain usage ↓ SSI
- Reduce OR personnel

Life moves us ➤

2012

Spine

LITERATURE REVIEW

A Methodological Systematic Review on Surgical Site Infections Following Spinal Surgery

Part 1: Risk Factors

Albert F. Pull ter Gunne, MD,* Allard J. F. Hoeman, MD, PhD,† David B. Cohen, MD, MPH,‡ Michael Schuetz, Dmed Habil, FRACS(Orth Surg),§ Cees J. H. M. van Laarhoven, MD, PhD,* and Joost J. van Middendorp, MD, PhD,§†¶

- 24 eligible studies
- 34 significant SSI factors identified
- **Diabetes, Obesity, Previous SSI most important**
 - Def: BMI >30
- Smoking only a risk factor in 2/11 studies investigating smoking

Life moves us ➤

Spine
LITERATURE REVIEW
SPINE Volume 37, Number 24, pp 2617-2631
©2012, Lippincott Williams & Wilkins

A Methodological Systematic Review on Surgical Site Infections Following Spinal Surgery

Part 1: Risk Factors

Albert F. Poller, MD, MPH, PhD,* Allard J. F. Hoeman, MD, PhD,† David B. Cohen, MD, MPH,‡
Michael Schuetz, Dmed Habil, FRACS(Orth Surg),§ Cees J. H. M. van Laarhoven, MD, PhD,*
and Joost J. van Middendorp, MD, PhD§§

- Prophylactic Strategies Identified:
 - Antibiotics
 - Skin antisepsis
 - Hair removal
 - Air ventilation system
 - Number of changes of OR personnel during surgery
 - Surgical clothing
 - Closed suction drains (if using a drain)

Life moves us ➤

2010

The Influence of Perioperative Risk Factors and Therapeutic Interventions on Infection Rates After Spine Surgery
A Systematic Review

James M. Schuster, MD, PhD,* Glenn Rachline, MD,† Daniel C. Norvell, PhD,‡
and Joseph R. Dettori, PhD†

- Evidence review 1990-2009
- ↑risks – age, obesity, diabetes, malnutrition, ASA, post appr, blood transfusion
- Unable to predict exact risk and unable to stratify

Table 7. List of Adjunct Measures for Preventing Postoperative Surgical-Site Infection Published in the Literature

Spine Surgery	General Orthopedic Surgery
Ultraclean air technology ²²	Ultraclean air technology ²⁵⁻²⁹
Betadine solution ^{20,22}	Betadine solution ^{23,25}
Closed wound suction drainage ^{18,19}	Occlusive garments ²⁵
No shaving ²²	Double gloving ^{23,21}
Silver impregnated dressings ⁷	Body exhaust suits ²⁹
	Ultraviolet lighting ³⁴
	Antiseptic scrubs ^{35,36}
	Nasal swabs ^{27,28}

Only irrigation with dilute betadine showed moderate support to reduce infection

Life moves us ➤

2004

Spine Update
Prevention of Postoperative Infection in Patients Undergoing Spinal Surgery

Erwin M. Brown, FRCPATH,* Ian K. Pople, FRCS,† John de Louvois, FRCPATH,‡
Alan Hedges, PhD,§ Roger Bayston, FRCPATH,¶ Stephen M. Eisenstein, FRCS,|| and
Peter Lane, FRCS|| for the British Society for Antimicrobial Chemotherapy Working Party on Neurosurgical Infections

- Prolonged op
 - Irrigate with saline +/- iodine regularly
 - Do not use abs in irrigation
- Minimise use of drains



Life moves us ➤

2009

Reduction of Postoperative Spinal Implant Infection Using Gentamicin Microspheres

Alec C. Stoll, MD, Ed Becker, MD, Steven C. Ludwig, MD, Daniel Galt, MD,
and Cornelis A. Poelstra, MD, PhD

- Rabbit model
- SSI 75% vs 38%
- Biomaterial infection (Ti rods) 58% vs 23%

Life moves us ➤

2008

National Collaborating Centre for Women's and Children's Health

Surgical site infection

prevention and treatment of surgical site infection

Clinical Guideline
October 2008
Formed to produce guidelines for the NICE by NICE

NCG74

Life moves us ➤

NICE 2008 NCG74

- At least 5% pts undergoing surgery develop SSI
- Perhaps some evidence now outdated
 - Esp for Ortho / Spine

Life moves us ➤

NICE Preop

- Patient prep – wash, clothes
- Hair removal – clippers
- Abs – timing, pharmacokinetics, rpt dosing
- Staff prep – clothing in / out th, jewellery, nails, polish

Do not use the following routinely to reduce the risk of surgical site infections:

- hair removal
- nasal decontamination with topical antimicrobial agents targeting *Staphylococcus aureus* **X**
- mechanical bowel preparation.

Life moves us ➤

The NEW ENGLAND JOURNAL of MEDICINE

ESTABLISHED IN 1912 JANUARY 7, 2010 VOL. 362 NO. 1

Preventing Surgical-Site Infections in Nasal Carriers of *Staphylococcus aureus*

Lonsdale G.M. Bode, M.D., Jan A.J.W. Kluytmans, M.D., Ph.D., Heiman F.L. Wertheim, M.D., Ph.D., Diana Bogaers, I.C.P., Christina M.J.E. Vandenbroucke-Grauls, M.D., Ph.D., Robert Roosendaal, Ph.D., Annet Troelstra, M.D., Ph.D., Adrienne T.A. Box, B.A.Sc., Andreas Voss, M.D., Ph.D., Ingeborg van der Tweel, Ph.D., Alex van Belkum, Ph.D., Henri A. Verbrugh, M.D., Ph.D., and Margreet C. Vos, M.D., Ph.D.


Life moves us ➤

NICE Intraop

- Brush / pick nails, scrub, sterile gowns, double glove, wash hands between ops
- Skin prep
- Maintain pt homeostasis – temp, O2, perfusion
- Use a wound dressing

Do not use the following to reduce the risk of surgical site infections:

- non-iodophor-impregnated incise drapes. If an incise drape is required, use an iodophor-impregnated drape unless the patient has an iodine allergy
- insulin routinely to optimise postoperative blood glucose levels in patients who do not have diabetes
- diathermy for surgical incision
- wound irrigation **X**
- intracavity lavage
- Intraoperative skin re-disinfection or topical cefotaxime in abdominal surgery.




THE NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE

Chlorhexidine-Alcohol versus Povidone-Iodine for Surgical-Site Antisepsis

Rabih O. Darouiche, M.D., Matthew J. Wall, Jr., M.D., Kamal M.F. Itani, M.D., Mary F. Otterson, M.D., Alexandra L. Webb, M.D., Matthew M. Carrick, M.D., Harold J. Miller, M.D., Samir S. Awad, M.D., Cynthia T. Crosby, B.S., Michael C. Mosier, Ph.D., Atef AlSharif, M.D., and David H. Berger, M.D.



Life moves us ➤

NICE Postop

- Aseptic tech for CoD
- Sterile saline up to 48hrs
- Tap water after 48hrs
- May shower safely after 48hrs (???)
- Refer TVN for wounds healing by secondary intention

Do not use the following to reduce the risk of surgical site infections:

- topical antimicrobial agents for surgical wounds that are healing by primary intention
- Eusol and gauze, or moist cotton gauze or mercuric antiseptic solutions for surgical wounds that are healing by secondary intention.

Life moves us ➤

Main Summary CDC Jan 2014

- Modifiable Risk Factors:
 - Ab Prophylaxis
 - Choice – local guidelines
 - Timing – 1 hour prior incision, stop within 24hrs
 - Dose based on BMI, procedure >3hrs, xs blood loss
 - Skin prep ineffective
 - Razors – use clippers
 - Excess OR traffic
 - Inadequate dressing – keep covered 24-48hrs
 - Poor glucose control - <200mg/dL or <11mmol/L
 - Colonization pre-existing organisms – Tx remote infections, nasal screening
 - Inadequate intraop oxygen levels – FiO2 50%
 - incl. normothermia

Life moves us ➤

2009

Intraoperative Fraction of Inspired Oxygen Is a Modifiable Risk Factor for Surgical Site Infection after Spinal Surgery

Lisa L. Mangos, M.D., M.P.H.,¹ Sara E. Cosgrove, M.D., M.S.,¹ Elizabeth A. Martinez, M.D., M.H.S.,¹ Margaret G. Turker, B.S.,¹ David B. Cohen, M.D., M.P.H.,² Trish M. Pate, M.D., M.Sc.¹

- 104 SSI vs 104 controls
- SSI associated with:
 - Prolonged procedure ($p < 0.001$)
 - ASA ≥ 3 ($p < 0.001$)
 - Lumbosacral ($p < 0.02$)
 - Posterior approach ($p < 0.02$)
 - Instrumentation ($p < 0.03$)
 - Obesity ($p < 0.01$)
 - Razor shaving ($p < 0.02$)
 - **FiO₂ < 50% ($p < 0.001$)**

At last we can blame the Anaesthetist!

Life moves us

RETRACTION

Longer Times >> More Damage

Gejo Spine 1999

- Soft tissue management:
 - Wound size, op duration, tissue damage, wound edges, oxygenation, blood supply

Life moves us

Respect the Skin Edges

Life moves us

Do The WHO

Surgical Safety Checklist

World Health Organization Patient Safety

Before induction of anaesthesia (with at least nurse and anaesthetist)	Before skin incision (with nurse, anaesthetist and surgeon)	Before patient leaves operating room (with nurse, anaesthetist and surgeon)
Has the patient confirmed his/her identity, life, procedure, and consent? Is the site marked? Is the anaesthesia machine and medication check complete? Is the pulse oximeter on the patient and functioning? Does the patient have a: Known allergy? Difficult airway or aspiration risk? Risk of >500ml blood loss (or >100ml in children)? Yes, and two IV/central access and fluids planned	Confirm all team members have introduced themselves by name and role. Confirm the patient's name, procedure, and where the incision will be made. Has antibiotic prophylaxis been given within the last 60 minutes? Anticipated Critical Events To Surgeon: What are the critical or non-routine steps? How long will the case take? What is the anticipated blood loss? To Anaesthetist: Are there any patient-specific concerns? To Nursing Team: Has safety (including indicator results) been confirmed? Are there equipment issues or any concerns? Is essential imaging displayed?	Nurse Verbally Confirms: The name of the procedure Completion of instrument, sponge and needle counts Specimen labelling (load specimen labels aloud, including patient name) Whether there are any equipment problems to be addressed To Surgeon, Anaesthetist and Nurse: What are the key concerns for recovery and management of this patient?

This checklist is not intended to be comprehensive. Additions and modifications to fit local practice are encouraged.

Revised 1/2009 © WHO, 2009

Life moves us

Surgical safety checklist

Items to be checked before, during and after the operation

Before induction of anaesthesia	Before skin incision	Before patient leaves operating room
SIGN IN PATIENT HAS CONFIRMED: • IDENTITY • SITE • PROCEDURE • CONSENT SITE MARKED/NOT APPLICABLE ANAESTHESIA SAFETY CHECK COMPLETED PULSE OXIMETER ON PATIENT AND FUNCTIONING DOES PATIENT HAVE A: KNOWN ALLERGY? NO YES DIFFICULT AIRWAY/ASPIRATION RISK? NO YES, AND EQUIPMENT/ASSISTANCE AVAILABLE RISK OF >500ML BLOOD LOSS (OR >100ML IN CHILDREN)? NO YES, AND ADEQUATE INTRAVENOUS ACCESS AND FLUIDS PLANNED	CHECK CONFIRM ALL TEAM MEMBERS HAVE INTRODUCED THEMSELVES BY NAME AND ROLE SURGEON, ANAESTHESIA PROFESSIONAL AND NURSE VERBALLY CONFIRM: • PATIENT • SITE • PROCEDURE ANTICIPATED CRITICAL EVENTS SURGEON REVIEWS: WHAT ARE THE CRITICAL OR UNEXPECTED STEPS, OPERATIVE DURATION, ANTICIPATED BLOOD LOSS? ANAESTHESIA TEAM REVIEWS: ARE THERE ANY PATIENT-SPECIFIC CONCERNS? NURSING TEAM REVIEWS: HAS STERILITY (INCLUDING INDICATOR RESULTS) BEEN CONFIRMED? ARE THERE EQUIPMENT ISSUES OR ANY CONCERNS? HAS ANTIBIOTIC PROPHYLAXIS BEEN GIVEN WITHIN THE LAST 60 MINUTES? YES NOT APPLICABLE ESSENTIAL IMAGING DISPLAYED? YES NOT APPLICABLE	SIGN OUT NURSE VERBALLY CONFIRMS WITH THE TEAM: • THE NAME OF THE PROCEDURE RECORDED • THAT INSTRUMENT, SPONGE AND NEEDLE COUNTS ARE CORRECT (OR NOT APPLICABLE) • HOW THE SPECIMEN IS LABELED (INCLUDING PATIENT NAME) • WHETHER THERE ARE ANY EQUIPMENT PROBLEMS TO BE ADDRESSED SURGEON, ANAESTHESIA PROFESSIONAL AND NURSE REVIEW THE KEY CONCERNS FOR RECOVERY AND MANAGEMENT OF THIS PATIENT

Life moves us

Cardiff Experience

- 4557 procedures over 5 years 2007-2012
- SSI 5% (223)
- 1/10 SSI required debridement (21)
- 1/50 required ROM / Replacement (5)

Is Aggressive Surgery Necessary for Acute Postoperative Deep Spinal Wound Infection?

Hsu-Shan Hong, MD,* Ming-Chau Chang, MD,* Chien-Lin Liu, MD, and Tain-Hsiung Chen, MD

SPINE, Volume 33, Number 22, pp 2473-2478 ©2008, Lippincott Williams & Wilkins

Life moves us

Common Spinal SSI Rates

72/3230 = 2.2% Original Article
J Chin Med Assoc
 2004;67:398-402

Postoperative Spinal Deep Wound Infection: A Six-year Review of 3230 Selective Procedures

36/1568 = 2.3%
Risk factors for deep surgical site infections after spinal fusion

J. J. P. Schimmel · P. P. Horsting · M. de Kleuver · G. Wonders · J. van Limbeek

752/24,774 = 3%
Risk Factors for Postoperative Spinal Wound Infections After Spinal Decompression and Fusion Surgeries

Anand Veeravagu, MD,*† Chirag G. Patel, MD,*† Shivanand P. Lad, MD, PhD,*† and Maxwell Boskey, MD,*†

132/3174 = 4%
The Presentation, Incidence, Etiology, and Treatment of Surgical Site Infections After Spinal Surgery

Albert F. Pull ter Gunne, MD,* Ahmed S. Mohamed, MD,* Richard L. Skolasky, ScD,*† Goss J. H. M. van Laarhoven, MD, PhD,*† and David B. Cohen, MD, MPH

48/1095 = 4.4%
Risk Factors for Infection After Spinal Surgery

Andrew Fang, MD, Serena S. Hu, MD, Nathan Endres, MD, and David S. Bradford, MD

Rate may depend on how closely data is collected and scrutinized

Life moves us ➤

Spine DEFORMITY

Rates of Infection After Spine Surgery Based on 108,419 Procedures

A Report from the Scoliosis Research Society Morbidity and Mortality Committee

Justin S. Smith, MD, PhD,* Christopher L. Shaffrey, MD,* Charles A. Sansur, MD,*† Sigurd H. Berven, MD,*† Kai-Ming C. Fu, MD, PhD,* Paul A. Broaddone, MD,*† Theodore J. Cherniack, MD,* Michael J. Goytan, MD,*† Hilali H. Noordeen, MD,*† Dennis R. Knapp, Jr., MD,*† Robert A. Hart, MD,*† William F. Donaldson III, MD,*† David W. Polly Jr, MD,*† Joseph H. Perrin, MD,*† Osheneba Boachie-Adjei, MD,*††

- SSI 2.1% (sup 0.8 deep 1.3)
- ↑revisions, fusions, implants
- Kyphosis > Scoliosis > Spondylolisthesis > Degen
 - Scoliosis and kyphosis - ↑neuromuscular (upto 14%)
- ↓MIS – discectomy and TLIF

Life moves us ➤

Spine DEFORMITY

Rates of Infection After Spine Surgery Based on 108,419 Procedures

A Report from the Scoliosis Research Society Morbidity and Mortality Committee

Surgical site infection rates after minimally invasive spinal surgery

Clinical article
 Jones E. O'Toole, MD,*† Krist M. Eckmiller, MD,*† and Richard G. Fessler, MD, PhD,*†

Infection Type	Approach	n	Rate (%)
Total infection (P=0.005)	Open	6,241	2.9%
	Minimally invasive	848	1.3%
Superficial infection (P=1.0)	Open	6,241	1.0%
	Minimally invasive	848	0.9%
Deep infection (P<0.001)	Open	6,241	1.9%
	Minimally invasive	848	0.4%

MIS: Less Pain Less Infection Less Bleeding Short stay

For TLIF and Discectomy

Life moves us ➤

Spine EPIDEMIOLOGY

Surgical Site Infections in Spine Surgery

Identification of Microbiologic and Surgical Characteristics in 239 Cases

Amir Abdul-jabbar, MD,* Sigurd H. Berven, MD,*† Serena S. Hu, MD,*† Dean Chou, MD,*† Praveen V. Mummaneni, MD,*† Steven Takemoto, PhD,*† Christopher R. Ames, MD,*† Dean Chou, MD,*† Bobby Jay, MD,*† Phil Weinstein, MD,*† Shane Burch, MD,*† and Catherine Liu, MD,*†

- 239 / 7529 cases SSI (3%)
 - Staph aureus 45%
 - Staph epi 31%
 - MRSA 34% and more common in revisions
 - Gram neg 30%
 - Sacral procedures – ↑gram neg and polymicrobia
 - Cervical procedures - ↓gram neg

Life moves us ➤

Spine SURGERY

6628 hospital visits SSI 2.9%

Surgical Site Infection in Spinal Surgery

Description of Surgical and Patient-Based Risk Factors for Postoperative Infection Using Administrative Claims Data

Coding study
 Amir Abdul-jabbar, BA, Steven Takemoto, PhD, Michael H. Weber, MD, PhD, FRCS, Serena S. Hu, MD, Praveen V. Mummaneni, MD, Vedat Deviren, MD, Christopher R. Ames, MD, Dean Chou, MD, Philip R. Weinstein, MD, Shane Burch, MD, and Sigurd H. Berven, MD

- Procedure:
 - Sacrum = 9.6%
 - Fuse > 7 = 7.8%
 - Fuse > 12 = 10.4%
 - Osteotomy = 6.5%
 - Op > 5hrs = 5.1%
 - Trans RBC = 5%
 - Trans Serum = 7.4%
 - Trans auto = 4.1%
- Patient:
 - Anaemia = 4.3%
 - Diabetes = 4.2%
 - Coronary art = 4.7%
 - Coagulopathy = 7.8%
 - Neoplasm = 5%

Life moves us ➤

Spinal Trauma

Early surgical site infections in adult spinal trauma: A prospective, multicentre study of infection rates and risk factors¹²

G. Lonjon^{1,2}, C. Dauzac³, E. Fourniols⁴, P. Guigui⁵, F. Bonnomet⁶, P. Bonneville⁶, The French Orthopaedic Surgery Traumatology Society⁷

Orthopaedics & Traumatology: Surgery & Research (2012) 98, 788–794

Risk Factors for Surgical Site Infection in the Patient With Spinal Injury

Oren G. Blam, MD,* Alexander R. Vaccaro, MD,*† Jed S. Vanichkachorn, MD,*† Todd J. Albert, MD,* Alan S. Hilibrand, MD,*† John M. Minnick, MD,*† and Sheila A. Murphy, MD†

- 6/169 (3.6%)
- 24/256 (9.4%)
- 7/54 (13%) controls vs 0/56 (0%) vanc in wound p=0.02

Life moves us ➤

2009

Surgical Site Infection in Spinal Metastasis
Risk Factors and Countermeasures

Satoru Demura, MD, Norio Kawahara, MD, Hidaki Murakami, MD, Koshi Nambu, MD, Satoshi Kato, MD, Katsuhito Yoshioka, MD, Tadaki Okuyama, MD, and Katsuro Tomita, MD

- Phase 1: 110 pts (113 ops)
 - 8/110 SSI (7.1%)
 - 7/22 SSI prior DXT (32%)
- Phase 2: 94 pts (97 ops)
 - Given PGE1 60micrograms bd post op mean 7/7
 - 3/97 SSI (3.1%)
 - 1/22 SSI prior DXR (4.5%)

NS

p<0.05

Life moves us


2011

Spine SURGERY

Postoperative Surgical Site Infections in Patients Undergoing Spinal Tumor Surgery
Incidence and Risk Factors

Ibrahim A. Omeis, MD, Mashaal Dhir, MD, Daniel M. Sciubba, MD, Oren N. Gottfried, MD, Matthew J. McGirt, MD, Frank J. Attenello, MD, Jean-Paul Wolinsky, MD, and Ziya L. Gokasan, MD

- 678 pts
- SSI:
 - Primary non bony = 8.9%
 - Metastatic = 9.5%
 - Primary bony = 13.7%



Life moves us

Paediatric Fusion

Risk Factors Associated With Surgical Site Infection After Pediatric Posterior Spinal Fusion Procedure

W. Matthew Linam, MD; Peter A. Margolis, MD, PhD; Mary Allen Staat, MD, MPH; Maria T. Britto, MD, MPH; Richard Hornung, DrPH; Amy Cassidy, PhD; Beverly L. Connolly, MD

Vol. 24 No. 8 INFECTION CONTROL AND HOSPITAL EPIDEMIOLOGY 591

SURGICAL-SITE INFECTION FOLLOWING SPINAL FUSION: A CASE-CONTROL STUDY IN A CHILDREN'S HOSPITAL

Annie-Claude Labbé, MD, FRCP(C); Anne-Marie Demers, MD, FRCP(C); Ramona Rodriguez, MSc; Vincent Arlet, MD; Kim Tangay, BSc; Dorothy L. Moore, PhD, MD, FRCP(C)

- 4.4%
- 5.2%

Life moves us

Paediatric Deformity SSI

Infection After Spinal Fusion for Pediatric Spinal Deformity
Thirty Years of Experience at a Single Institution

Patrick J. Cahill, MD,* Drew E. Warrick, MD,† Michael J. Lee, MD,‡ John Gaughan, PhD,§ Lawrence E. Vogel, MD,¶ Kim W. Hammerberg, MD,|| and Peter F. Sturm, MD,||

Diagnosis	No. Patients	No. Infections	Rate
Cerebral palsy	107	13	12.1
Myelomeningocele	84	15	17.9
Myopathies	27	1	2.7
SCI	37	6	16.2
Polio	12	4	33.0
Others	30	1	3.3

	No. Cases	No. Infections	Infection Rate (%)
Idiopathic	948	5	0.5
Congenital	139	2	1.4
Syndromic	166	5	3.0
Neuromuscular	323	43	13.3
Other	11	2	18.2

Deep Wound Infections After Neuromuscular Scoliosis Surgery
A Multicenter Study of Risk Factors and Treatment Outcomes

Paul D. Sponseller, MD, Dawn M. LaPorte, MD, Marc W. Hungerford, MD, Kevin Eck, MD, Keith H. Bridwell, MD, and Lawrence G. Lenke, MD

- 57 / 1543 = 3.7%
- 25 / 210 = 12%

Life moves us

Adult Deformity

Incidence of surgical site infection following adult spinal deformity surgery: an analysis of patient risk

Albert F. Poller Gunne · C. J. H. M. van Laarhoven · David B. Cohen


Eur Spine J (2010) 19:982–988
DOI 10.1007/s00586-009-1269-1

ORIGINAL ARTICLE






- 46/830 = 5.5%
- Risk – obesity and previous SSI


Life moves us

L4/5 Stenosis Facet Cyst



Life moves us





Risk Factors for Surgical Site Infection Following Orthopaedic Spinal Operations

By Margaret A. Olsen, PhD, MPH, Jeffrey J. Nepple, MD, K. Daniel Riew, MD, Lawrence G. Lenke, MD, Keith H. Bridwell, MD, Jennie Mayfield, BSN, MPH, CIC, and Victoria J. Fraser, MD

- 46/2316 ops -2% SSI
— vs 227 random controls
- Pre and within 5/7 post op Glucose
- Diabetes
- Poor timing of Abs
- Obesity
- 2 or more residents scrubbed

At last we can blame the Fellow!

Life moves us

Some Rules About...

When a Reg / Fellow starts:

1. Don't let them put their dirty hands in the wound
2. Don't let them do the operation
3. Don't let them close the wound
4. Supervise them closely!


Life moves us


Take Home Message

MRSA

Avoid operating on obese diabetics with previous history of spinal infection

Life moves us





Spine
LITERATURE REVIEW

SPINE Volume 37, Number 24, pp 2015-2019
©2012, Lippincott Williams & Wilkins

A Methodological Systematic Review on Surgical Site Infections Following Spinal Surgery

Part 1: Risk Factors

Albert F. Pull ter Gunne, MD,* Allard J. F. Hoosman, MD, PhD,† David. B. Cohen, MD, MPH,‡
Michael Schuetz, Dmed Habil, FRACS(Orth.Surg),§ Cees J. H. M. van Laarhoven, MD, PhD,*
and Joost J. van Middendorp, MD, PhD§¶

- **Patient:**
 - Age, sex, race, workers comp, height, prev spinal op, prev non spinal op, prev op (NOS), prev spinal infection / SSI, smoking, alcohol, immunosuppressant, obesity, diabetes, rheumatoid, cardiovasc dis, pulm dis, UTI, urinary incontinence, complete neuro deficit, functional status, Karnofsky score, ASA, NNIS risk index, malignancy, neoadjuvant therapy, tumour size, weight loss

Life moves us ➤

Spine
LITERATURE REVIEW

SPINE Volume 37, Number 24, pp 2015-2019
©2012, Lippincott Williams & Wilkins

A Methodological Systematic Review on Surgical Site Infections Following Spinal Surgery

Part 1: Risk Factors

Albert F. Pull ter Gunne, MD,* Allard J. F. Hoosman, MD, PhD,† David. B. Cohen, MD, MPH,‡
Michael Schuetz, Dmed Habil, FRACS(Orth.Surg),§ Cees J. H. M. van Laarhoven, MD, PhD,*
and Joost J. van Middendorp, MD, PhD§¶

- **Anaesthetic:**
 - Propofol, vent and oxygenation characteristics, body temp, intraop fluid infusion rate, clotting agents
- **Surgical:**
 - Indication, date, room personnel, specific OR, timing, hair removal, antisepsis, antibiotics, anatomical approach, type, number surgeries, duration, EBL, level of surgery, no of levels, graft type, dural tear, microscope, closed suction drain

Life moves us ➤

Spine
LITERATURE REVIEW

SPINE Volume 37, Number 24, pp 2015-2019
©2012, Lippincott Williams & Wilkins

A Methodological Systematic Review on Surgical Site Infections Following Spinal Surgery

Part 1: Risk Factors

Albert F. Pull ter Gunne, MD,* Allard J. F. Hoosman, MD, PhD,† David. B. Cohen, MD, MPH,‡
Michael Schuetz, Dmed Habil, FRACS(Orth.Surg),§ Cees J. H. M. van Laarhoven, MD, PhD,*
and Joost J. van Middendorp, MD, PhD§¶

- **Postoperative:**
 - Steroids, bed bath, post op incontinence, any incontinence
- **Hospitalization:**
 - Preop, post op location, LOS

Life moves us ➤

Spine
LITERATURE REVIEW

SPINE Volume 37, Number 24, pp 2015-2019
©2012, Lippincott Williams & Wilkins

A Methodological Systematic Review on Surgical Site Infections Following Spinal Surgery

Part 1: Risk Factors

Albert F. Pull ter Gunne, MD,* Allard J. F. Hoosman, MD, PhD,† David. B. Cohen, MD, MPH,‡
Michael Schuetz, Dmed Habil, FRACS(Orth.Surg),§ Cees J. H. M. van Laarhoven, MD, PhD,*
and Joost J. van Middendorp, MD, PhD§¶

- **Laboratory:**
 - Preop: malnutrition, albumin, protein, WCC, Cr, Hct, Glu
 - Post op: Glu, Hb, Transfusion – pRBC after op, FFP intraop

Life moves us ➤

Spine
LITERATURE REVIEW

SPINE Volume 37, Number 24, pp 2015-2019
©2012, Lippincott Williams & Wilkins

A Methodological Systematic Review on Surgical Site Infections Following Spinal Surgery

Part 2: Prophylactic Treatments

Joost J. van Middendorp, MD, PhD,†† Albert F. Pull ter Gunne, MD,§
Michael Schuetz, Dmed Habil, FRACS(Orth.Surg),†† David B. Cohen, MD, MPH,‡
Allard J. F. Hoosman, MD, PhD,† and Cees J. H. M. van Laarhoven, MD, PhD§

- **18 eligible studies**
- **Main Reported Prophylactic Strategies in Lit:**
 - Antibiotics, closed suction wound drains, laminar flow, body exhaust suits, routine UV lights in theatre, povidone-iodine wound irrigation, Ab wound irrigation, chlorhex shower night before op, frequent dressing changes, debridement non viable tissue intraop, shaving preop, silverlon dressings, use of PGE1

Life moves us ➤

Cardiff Ab Prophylaxis 2014

- Non Instrumented Low Risk:
 - Cefuroxime 1.5g Induction
 - Allergy: Teicoplanin 400mg + Gentamicin 3mg/Kg
- High Risk and Instrumented:
 - Cefuroxime 1.5g and Teicoplanin 400mg Induction
 - Post Op Cef 750mg x2-3 8° and Teic 400mg x1 @12hrs
 - Allergy: Teicoplanin 400mg + Gentamicin 3mg/Kg
 - Post Op Teic 400mg x1 @12hrs
- If post op catheter insertion:
 - 160mg Gentamicin

Risk Factors for Surgical Site Infection Following Orthopaedic Spinal Operations

By Margaret A. Chen, PhD, MPH, Jeffrey I. Noppa, MD, E. David Rowe, MD, Lawrence G. Jenkins, MD, Keith H. Baidoo, MD, Suresh Mayhew, MD, MPH, CJC, and Vinita I. Frank, MD

TABLE II Univariate Comparisons of Individual Risk Factors in Patients with and without Surgical Site Infection Following Orthopaedic Spinal Operations

Patient-Level Characteristics	No. (%) of Patients with Surgical Site Infection (N = 46)	No. (%) of Uninfected Patients (N = 227)	Odds Ratio (95% Confidence Interval)	P Value
Body mass index				
Normal (≤ 25 kg/m ²)	9 (20)	84 (37)	1.0	
Overweight (>25 and <30 kg/m ²)	11 (24)	77 (34)	1.3 (0.5, 3.4)	0.546
Obese (≥ 30 and ≤ 35 kg/m ²)	18 (39)	37 (16)	4.5 (1.9, 11.0)	0.001
Morbidly obese (>35 kg/m ²)	8 (17)	29 (13)	2.6 (0.9, 7.3)	0.075
Diabetes and hyperglycemia				
Neither (no diabetes or hyperglycemia)	20 (43)	179 (79)	1.0	
Serum glucose >75 th percentile	12 (26)	33 (15)	3.3 (1.5, 7.3)	0.004
Diabetes	14 (30)	15 (7)	8.4 (3.5, 19.8)	<0.001
Diagnoses				
Herniated disc	6 (13)	63 (28)	0.4 (0.2, 1.0)	0.036
Fracture	6 (13)	9 (4)	3.6 (1.2, 10.8)	0.025
American Society of Anesthesiologists class 3 or 4	22 (48)	59 (26)	2.6 (1.4, 5.0)	0.003
Nerve root compression	22 (48)	153 (67)	0.4 (0.2, 0.8)	0.012
Transfusion				
Cell Saver or autologous blood	10 (22)	35 (15)	2.2 (0.9, 5.0)	0.071
Packed red blood cells or platelets	15 (33)	33 (14.5)	3.4 (1.6, 7.4)	0.001
Postoperative incontinence	8 (17)	14 (6)	3.2 (1.3, 8.2)	0.018
Preoperative or postoperative incontinence	13 (28)	27 (12)	2.9 (1.4, 6.2)	0.004

Risk Factors for Surgical Site Infection Following Orthopaedic Spinal Operations

By Margaret A. Chen, PhD, MPH, Jeffrey I. Noppa, MD, E. David Rowe, MD, Lawrence G. Jenkins, MD, Keith H. Baidoo, MD, Suresh Mayhew, MD, MPH, CJC, and Vinita I. Frank, MD

TABLE III Univariate Comparisons of Surgical Risk Factors in Patients with and without Surgical Site Infection Following Orthopaedic Spinal Operations

Operative Characteristics	No. (%) of Patients with Surgical Site Infection (N = 46)	No. (%) of Uninfected Patients (N = 227)	Odds Ratio (95% Confidence Interval)	P Value
Cervical level	11 (24)	97 (43)	0.4 (0.2, 0.9)	0.017
Posterior approach	42 (91)	172 (76)	3.4 (1.2, 9.8)	0.020
Suboptimal timing of prophylactic antibiotic therapy*	15 (33)	31 (14)	3.1 (1.5, 6.3)	0.002
Suboptimal dosage of prophylactic antibiotic†	20 (43.5)	51 (22.5)	2.7 (1.4, 5.1)	0.003
Only cefazolin used as prophylactic antibiotic	24 (52)	168 (74)	0.4 (0.2, 0.7)	0.003
Aminoglycoside used as prophylactic antibiotic	17 (37)	41 (18)	2.7 (1.3, 5.3)	0.004
Intravenous steroids during operation	8 (17)	85 (37)	0.4 (0.2, 0.8)	0.009
Wound irrigated with antibiotic-containing solution	40 (87)	162 (71)	2.7 (1.1, 6.6)	0.028
No. of intervertebral levels				
1	10 (22)	83 (37)	1.0	
2-3	21 (46)	94 (41)	1.9 (0.8, 4.2)	0.135
4-6	7 (15)	30 (13)	1.9 (0.7, 5.5)	0.218
≥ 7	8 (17)	20 (9)	3.3 (1.2, 9.5)	0.025
Duration of operation >75 th percentile‡	18 (39)	49 (22)	2.4 (1.2, 4.6)	0.012
Hemovac drain placed	26 (56.5)	95 (42)	1.8 (1.0, 3.4)	0.068
Drains in place ≥ 3 days	33 (72)	108 (48)	2.8 (1.4, 5.6)	0.003
≥ 2 resident surgeons	32 (70)	109 (48)	2.5 (1.3, 4.9)	0.008

*Cefazolin given more than sixty minutes before the incision or after the incision, or another antibiotic given after the incision. Two hundred and twenty-nine (84%) of the 273 patients received prophylactic cefazolin, alone (192), in combination with an aminoglycoside (thirty-four), or in combination with another antibiotic (three). †One gram of cefazolin used as a prophylactic antibiotic in patients with a body mass index of ≥ 30.0 kg/m². ‡The 75th percentile for fusion was 310.5 minutes; the 75th percentile for non-fusion operations was 145 minutes.

Postoperative Wound Infection Rates after Posterior Instrumented Spinal Surgery in Diabetic Patients

Jen-Chung Liao, MD; Wen-Jer Chen, MD; Lih-Huei Chen, MD; Chi-Chien Niu, MD

- Diabetes
- 337 Posterior lumbar fusions
- 39 diabetics – 4 had SSI
- 298 non diabetic – 2 had SSI

TABLE I. Rate of Infection Based on Patient Age and Primary Diagnosis*

Primary Diagnosis	No. Cases	Superficial Infection (%)	Deep Infection (%)	Total Infection (%)
Adult (≥ 21 yr)				
Degenerative disease	46,434	308 (0.7)	342 (0.7)	650 (1.4)
Scoliosis	5801	66 (1.1)	146 (2.5)	212 (3.7)
Spondylolisthesis	10,529	93 (0.9)	130 (1.2)	223 (2.1)
Fracture	6025	31 (0.5)	92 (1.5)	123 (2.0)
Kyphosis	2012	28 (1.4)	57 (2.8)	85 (4.2)
Other	11,089	113 (1.0)	237 (2.1)	350 (3.2)
Primary Tumor†	404	4 (1.0)	7 (1.7)	11 (2.7)
Metastatic Tumor†	726	9 (1.2)	27 (3.7)	36 (5.0)
Acute osteomyelitis†	866	12 (1.4)	32 (3.7)	44 (5.1)
Implant revision/removal†	1720	16 (0.9)	39 (2.3)	55 (3.2)
Not recorded	192	2 (1.0)	2 (1.0)	4 (2.1)
Total	82,082	641 (0.8)	1006 (1.2)	1647 (2.0)
Pediatric (<21 yr)				
Degenerative disease	654	5 (0.8)	1 (0.2)	6 (0.9)
Scoliosis	20,424	195 (1.0)	341 (1.7)	536 (2.6)
Spondylolisthesis	827	9 (1.1)	12 (1.5)	21 (2.5)
Fracture	623	5 (0.8)	9 (1.4)	14 (2.2)
Kyphosis	1555	35 (2.3)	49 (3.2)	84 (5.4)
Other	1273	11 (0.9)	23 (1.8)	34 (2.7)
Primary tumor	152	0 (0.0)	5 (3.3)	5 (3.3)
Implant revision/removal†	308	1 (0.3)	6 (1.9)	7 (2.3)
Not recorded	76	1 (1.3)	1 (1.3)	2 (2.6)
Total	25,432	261 (1.0)	436 (1.7)	697 (2.7)

*Age not recorded for 905 patients. These patients are not included in this table.

†Selected subgroup within the "other" category. Note that numbers of cases for these subgroups are part of the total other cases indicated.

TABLE 2. Rate of infection among adult patients with a primary diagnosis of degenerative disease, stratified based on spinal location and subtype of disease

Spinal Region/Diagnosis	No. Cases	Superficial Infection (%)	Deep Infection (%)	Total Infection (%)
Cervical				
Postlaminectomy syndrome	35	1 (2.9)	1 (2.9)	2 (5.7)
Spinal stenosis	3466	13 (0.4)	27 (0.8)	40 (1.2)
Spondylotic radiculopathy	1859	9 (0.5)	5 (0.3)	14 (0.8)
Degenerative disc disease	1769	5 (0.3)	7 (0.4)	12 (0.7)
Disc herniation	4472	11 (0.2)	12 (0.3)	23 (0.5)
Not recorded	73	0 (0.0)	0 (0.0)	0 (0.0)
Total	11,674	39 (0.3)	52 (0.4)	91 (0.8)
Thoracic				
Degenerative disc disease	95	0 (0.0)	3 (3.2)	3 (3.2)
Spinal stenosis	161	2 (1.2)	2 (1.2)	4 (2.5)
Disc herniation	235	3 (1.3)	1 (0.4)	4 (1.7)
Spondylotic radiculopathy	17	0 (0.0)	0 (0.0)	0 (0.0)
Not recorded	16	0 (0.0)	0 (0.0)	0 (0.0)
Total	524	5 (1.0)	6 (1.1)	11 (2.1)
Lumbar				
Spondylotic radiculopathy	949	22 (2.3)	14 (1.5)	36 (3.8)
Spinal stenosis	12,270	116 (0.9)	153 (1.2)	269 (2.2)
Postlaminectomy syndrome	573	6 (1.0)	4 (0.7)	10 (1.7)
Degenerative disc disease	7213	55 (0.8)	58 (0.8)	113 (1.6)
Disc herniation	12,694	62 (0.5)	51 (0.4)	113 (0.9)
Not recorded	211	2 (0.9)	2 (0.9)	4 (1.9)
Total	33,910	263 (0.8)	282 (0.8)	545 (1.6)
Spinal location not recorded	326	1 (0.3)	2 (0.6)	3 (0.9)



TABLE 3. Rate of Infection Among Patients with a Primary Diagnosis of Scoliosis, Stratified Based on Patient Age and Subtype of Scoliosis

Age Group/Type of Scoliosis	No. of Cases	Superficial Infection (%)	Deep Infection (%)	Total Infection (%)
Adult (≥21 yr)				
Neuromuscular	292	8 (2.7)	18 (6.2)	26 (8.9)
Posttraumatic	30	0 (0.0)	2 (6.7)	2 (6.7)
Degenerative	2533	31 (1.2)	73 (2.9)	104 (4.1)
Congenital	137	1 (0.7)	4 (2.9)	5 (3.6)
Idiopathic	2488	23 (0.9)	46 (1.8)	69 (2.8)
Other	139	3 (2.2)	2 (1.4)	5 (3.6)
Not recorded	182	0 (0.0)	1 (0.5)	1 (0.5)
Total	5801	66 (1.1)	146 (2.5)	212 (3.7)
Pediatric (<21 yr)				
Posttraumatic	35	1 (2.9)	1 (2.9)	2 (5.7)
Neuromuscular	4855	83 (1.7)	184 (3.8)	267 (5.5)
Congenital	2045	27 (1.3)	18 (0.9)	45 (2.2)
Idiopathic	11,741	65 (0.6)	100 (0.9)	165 (1.4)
Other	1464	16 (1.1)	31 (2.1)	47 (3.2)
Not recorded	284	3 (1.1)	7 (2.5)	10 (3.5)
Total	20,424	195 (1.0)	341 (1.7)	536 (2.6)

Life moves us

TABLE 4. Rate of Infection Among Patients with a Primary Diagnosis of Kyphosis, Stratified Based on Patient Age and Subtype of Kyphosis

Age Group/Type of Kyphosis	No. Cases	Superficial Infection (%)	Deep Infection (%)	Total Infection (%)
Adult (≥21 yr)				
Postlaminectomy	198	5 (2.5)	5 (2.5)	10 (5.1)
Osteoporosis	104	3 (2.9)	2 (1.9)	5 (4.8)
Scheuermann	227	2 (0.9)	8 (3.5)	10 (4.4)
Posttraumatic	365	3 (0.8)	12 (3.3)	15 (4.1)
Fixed sagittal plane deformity	553	6 (1.1)	15 (2.7)	21 (3.8)
Neuromuscular	42	0 (0.0)	1 (2.4)	1 (2.4)
Congenital	48	0 (0.0)	1 (2.1)	1 (2.1)
Other	396	9 (2.3)	13 (3.3)	22 (5.6)
Not recorded	79	0 (0.0)	0 (0.0)	0 (0.0)
Total	2012	28 (1.4)	57 (2.8)	85 (4.2)
Pediatric (<21 yr)				
Neuromuscular	264	10 (3.8)	27 (10.2)	37 (14.0)
Fixed sagittal plane deformity	18	1 (5.6)	0 (0.0)	1 (5.6)
Postlaminectomy	64	3 (4.7)	0 (0.0)	3 (4.7)
Scheuermann	634	12 (1.9)	15 (2.4)	27 (4.3)
Posttraumatic	31	1 (3.2)	0 (0.0)	1 (3.2)
Congenital	329	3 (0.9)	2 (0.6)	5 (1.5)
Other	200	4 (2.0)	5 (2.5)	9 (4.5)
Not recorded	15	1 (6.7)	0 (0.0)	1 (6.7)
Total	1555	35 (2.3)	49 (3.2)	84 (5.4)

Life moves us

TABLE 5. Summary of Surgical Characteristics Assessed for Association with Infection

Parameter	No. Cases	Superficial Infection (%)	Deep Infection (%)	Total Infection (%)	P*
Spinal fusion performed					
Yes	72,534	642 (0.9)	1072 (1.5)	1714 (2.4)	<0.001
No	35,877	267 (0.7)	377 (1.1)	644 (1.8)	
Not recorded	8	0 (0.0)	0 (0.0)	0 (0.0)	
Type of fusion					
Post-ant-post	271	2 (0.7)	7 (2.6)	9 (3.3)	
Anterior-posterior	7887	91 (1.2)	161 (2.0)	252 (3.2)	<0.001†
Posterolateral	19,710	221 (1.1)	379 (1.9)	600 (3.0)	<0.001†
Interlaminar/facet	16,192	167 (1.0)	281 (1.7)	448 (2.8)	0.003†
All fusion cases	72,534	642 (0.9)	1072 (1.5)	1714 (2.4)	
TLIF/PLIF	12,267	104 (0.8)	182 (1.5)	286 (2.3)	
Anterior only	15,336	46 (0.3)	48 (0.3)	94 (0.6)	<0.001†
Not recorded	871	10 (1.1)	14 (1.6)	24 (2.8)	
Implants					
Yes	74,114	647 (0.9)	1092 (1.5)	1739 (2.3)	<0.001
No	34,305	262 (0.8)	357 (1.0)	619 (1.8)	
Revision surgery					
Yes	16,503	183 (1.1)	359 (2.2)	542 (3.3)	<0.001
No	91,916	726 (0.8)	1090 (1.2)	1816 (2.0)	
Minimally invasive approach					
Yes	14,301	43 (0.3)	35 (0.2)	78 (0.5)	<0.001
No	94,115	866 (0.9)	1414 (1.5)	2280 (2.4)	
Not recorded	3	0 (0.0)	0 (0.0)	0 (0.0)	

*P-values are based on statistical comparisons of total infection rates.

†Anterior-posterior, posterolateral, and interlaminar/facet fusions had significantly higher and anterior only had significantly lower rates of infection compared with the overall rate of infection for all fusion patients (2.4%).

TLIF indicates transforaminal lumbar interbody fusion; PLIF, posterior lumbar interbody fusion.

Life moves us